



CAID	Coyle-Ashport frequently flooded-Ironmound complex, 1 to 8 percent slopes	17.33	5.5%	IVe		4713	5	5	12	1	1		3	1	2
GraC	Grainola silty clay loam, 3 to 5 percent slopes	10.58	3.3%	IIIe		3724	2	3	18	1	1				
ColC2	Coyle-Ironmound complex, 3 to 5 percent slopes, eroded	9.15	2.9%	IIIe		0	3	3	14				3	1	2
W	Water	5.94	1.9%	VIII		0									
RnnC2	Renthin silty clay loam, 3 to 5 percent slopes, eroded	5.66	1.8%	IIIe		0	4	5	19		28				
PieB	Piedmont silty clay loam, 1 to 3 percent slopes	3.87	1.2%	IIIs		3849	4	5	24				4	2	
ZaHC	Zaneis-huska complex, 3 to 5 percent slopes	2.76	0.9%	IIIe		4126	4	4	19	1	1		3	1	3
KrkB	Kirkland silty clay loam, 1 to 3 percent slopes	1.90	0.6%	IIIe		3853	4	5	29		1		4	2	4
<b>Weighted Average</b>						<b>2479.6</b>	<b>3.7</b>	<b>3.7</b>	<b>19.6</b>	<b>3.6</b>	<b>5.1</b>	<b>0.1</b>	<b>1.1</b>	<b>0.3</b>	<b>0.2</b>

\*n: The aggregation method is "Weighted Average using major components"

\*c: Using Capabilities Class Dominant Condition Aggregation Method

Soils data provided by USDA and NRCS.